arXiv:hep-ph/9312234v1 6 Dec 1993

Cancellation of power enhancements in non-spectator decays

U. Aglietti
SISSA-ISAS, Via Beirut 2, 34014 Trieste, Italy
INFN, Sezione di Trieste, Via Valerio 2, 34100 Trieste, Italy

Abstract

Exclusive non-spectator decay rates of beauty hadrons contain power enhancements of the form $(m_b/m)^2$ and m_b/m, where m_b is the b-quark mass and m is a light quark mass. An implicit argument has been recently given according to which these singularities cancel in the totally inclusive decay width. We present in this note a completely explicit computation of the diagrams containing power enhancements. Our results agree with the previous general conclusion.

The basic decay mechanism of beauty hadrons is the decay of the constituent b-quark. The spectator model assumes the dominance of this mechanism, neglects binding corrections of the b-quark in the hadron, and computes inclusive decay widths by means of the parton model. For example, the semileptonic decay $H_b \ \rightarrow \ X_c + l + \nu$, where H_b is a beauty hadron and X_c is any hadronic final state with charm $C = 1$, is computed as the decay $b \rightarrow c + l + \nu$. The rate is given by:

$$\Gamma \ = \ \Gamma_0[1 - (2\alpha_s/3\pi) f(m_c/m_b)] \ I(m_c/m_b, m_l/m_b, 0) \tag{1}$$

where $\Gamma_0 = G_F^2 m_b^5 \mid V_{cb} \mid^2 /192\pi^3$, $f(x)$ is a perturbative QCD correction with $f(0) = \pi^2 - 25/4$, and $I(x_1, x_2, x_3)$ is a phase-space correction factor for non-zero masses in the final state, with $I(0, 0, 0) = 1$ [1].
Hadronic decays are computed in a quite analogous way. The rate for the decay

$$b \ \rightarrow \ c + \overline{u} + d, \tag{2}$$

for example, is given by:

$$\Gamma = 3\Gamma_0 \mid V_{ud} \mid^2 \frac{2C_+^2 + C_-^2}{3} J I(m_c/m_b, m_d/m_b, m_u/m_b) \tag{3}$$

where $C_{+,-} = (\alpha_s(m_b)/\alpha_s(m_W))^{d+,-}$, with $d_{+,-} = -6/23, \ 12/23$, are coefficients resumming leading-logs coming from the renormalization of the non-leptonic hamiltonian [2] (numerically, $C_+ \simeq 0.88$ and $C_- \simeq 1.29$), and J is a factor including QCD-subleading corrections computed in ref.[3].
An important consequence of the spectator model is that all beauty hadrons should have the same lifetime, because the light quark(s) in the initial state behave like passive spectators. LEP experiments aim at testing this prediction with errors less than the expected splitting. The semileptonic branching fraction B_{SL} can also be predicted with reasonable accuracy. This is because the leading dependence on m_b^5 (m_b is not clearly defined), cancels in taking the ratio of widths. A detailed computation including also penguin diagrams, gives a semileptonic branching-ratio in the range [4]: $B_{SL} \ = \ [12.2 \pm 0.45(scale) \pm 0.8(\alpha_s)]\%$, for current values of quark masses, and $B_{SL} \ = \ [14.4 \pm 0.45(scale) \pm 0.8(\alpha_s)]\%$, for constituent values of quark masses. The first uncertainty comes from the scale of the coupling α_s, while the second one comes from the experimental error on $\alpha_s(m_Z)$. The experimental value $B_{SL} \ = \ (10.5 \pm 0.5)\%$[5] is marginally consistent with the

prediction of the spectator model, implying non-spectator effects of the order of 15%.

There is a clear indication of the relevance of decay mechanisms which lie outside the spectator model, involving the light quark(s) in the hadron in an essential way (non-spectator mechanisms).

Non-spectator mechanisms are of various kind and involve different quark-level processes. The corresponding rates are proportional to m_b^n with $n \leq 3$. Since the partonic rate is proportional to m_b^5, all non-spectator effects are preasymptotic in the limit of a very large mass of the heavy quark, $m_b \to \infty$.

Let us recall very briefly the corrections related to the (not so free) b-decay. Partonic rates, such as those in eqs.(1) and (3), have corrections for the binding of the b-quark in the hadron (bound-state corrections [6]), as well as non-perturbative corrections for final state interactions. Recently there has been some progress in the model-independent computation of this kind of corrections using the heavy quark effective theory ($HQET$) [7]. The semileptonic branching fraction of B mesons is modified according to [8]:

$$\frac{\delta B_{SL}(B)}{B_{SL}(B)} \approx 6 \frac{M_{B*}^2 - M_B^2}{m_b^2} \frac{C_+^2 - C_-^2}{2N} \approx -2\% \tag{4}$$

B_{SL} therefore is reduced with respect to the spectator value in the 'right direction', although by a tiny amount only. The effect is small because color dynamics gives a large suppression factor: $(C_+^2 - C_-^2)/2N \simeq -0.15$. $\delta B_{SL}/B_{SL}$ vanishes instead in the Λ_b case. As it stems from eq.(4), these corrections are of order $1/m_b^2$ with respect to the partonic rate.

We now consider corrections related to different quark-level processes with respect to b decay. The case of B-mesons is discussed for simplicity. The neutral B_d meson can decay via annihilation of the constituents b and $\overline{d}$ quarks [9]. The corresponding decay rate is given by:

$$\frac{\delta\Gamma_{ann}}{\Gamma_0} = 24\pi^2 \left(\frac{f_B}{m_b}\right)^2 \left(\frac{m_c}{m_b}\right)^2 \left[3\left(\frac{C_+ - C_-}{2}\right)^2 + \frac{C_+^2 + C_-^2}{2} + \frac{1}{3}\left(\frac{C_+ + C_-}{2}\right)^2\right] \tag{5}$$

where f_B is the decay constant of the B meson (numerically, $f_B \sim 200\ MeV$ from lattice QCD computations [10]). The width in eq.(5) is very small, $\delta\Gamma_{ann}/\Gamma_0 \sim 5\%$, because there is a large helicity-suppression factor $(m_c/m_b)^2 \sim 0.1$. The annihilation mechanism increases the decay rate of the B^0 meson with respect to the spectator value.

The decay of charged B_u mesons is modified with respect to the spectator case by an interference effect, because in the channel (2) there are two identical $\overline{u}$-quarks in the final state [11, 12]. The modification of the decay rate due to the interference is given by:

$$\frac{\delta\Gamma_{int}}{\Gamma_0} = 48\pi^2 \left(\frac{f_B}{m_b}\right)^2 \left[\frac{C_+^2 - C_-^2}{2} + \frac{C_+^2 + C_-^2}{2N}\right] \tag{6}$$

The rate in eq.(6) is negative, i.e. the interference reduces the decay rate of the charged B's with respect to the partonic value. In this case, there is no more the helicity-suppression factor present in eq.(5), but the non-spectator rate is again very small because of color dynamics. There is a large cancellation between the two terms in the square bracket in eq.(6), which is efficient up to $\sim 90\%$, and gives $\delta\Gamma_{int}/\Gamma_0 \sim -3\%$.

As it stems from eqs.(5) and (6), the annihilation and interference rates are proportional to m_b^2, because the decay constant scales as $f_M \propto 1/\sqrt{m_b}$ for very large masses [13]. These mechanisms therefore are subleading with respect to the non-perturbative corrections discussed before. As a consequence, we expect for very large m_b a greater difference in lifetimes between mesons and baryons, than the ones inside mesons.

Radiative corrections to non-spectator decays may in principle increase considerably the rates. In the case of the interference, for example, gluon radiation from the constituent light quark contributes to the decay rate by a term of the form:

$$\frac{\delta\Gamma_{int}'}{\Gamma_0} = \pi\alpha_s \left(\frac{f_B}{m}\right)^2 (1 - 1/N^2)\frac{C_+^2 + C_-^2}{2} \tag{7}$$

There is a power enhancement of the form $(m_b/m)^2$ with respect to the lowest order term, which overcomes the typical suppression factor $\alpha_s/4\pi$ of the radiative corrections. Furthermore, color dynamics is different from that one of the tree amplitude, and the almost complete cancellation between C_+ and C_- terms in eq.(6) does not occur any more. Formula (7) implies that non-spectator effects may be substantially greater than 1% level as implied by the tree-level computations (5) and (6). However, the rate in eq.(7) depends critically on the value of the light quark mass m, which cannot be reliably estimated since it involves unknown non-perturbative dynamics. The result therefore has a limited phenomenological impact. Analogous power enhancements occur in the radiative corrections to the annihilation width of B^0 mesons.

In ref.[14] a brilliant analyticity argument has been given according to which all power enhancements cancel in the totally inclusive decay rate, leaving only residual logarithmic singularities of the form $\log(m_b/m)$.

In this note we present an explicit computation of the diagrams containing power enhancements. For definiteness we consider the interference case. The absorptive part of the diagram in fig.1 is given by the sum of the cuts (i), (ii) and (iii):

$$\begin{aligned}
\mathcal{M} \;=\; & \mathcal{M}_{(i)} + \mathcal{M}_{(ii)} + \mathcal{M}_{(iii)} = C \int\!\!\int \frac{d^4k}{(2\pi)^4} \frac{d^4l}{(2\pi)^4} \overline{v}_u(p_2) i g \gamma_\sigma \\
& i \frac{-\hat{k} - \hat{p}_2 + m}{(k-p_2)^2 - m^2 + i\epsilon} (-i) \frac{G_F}{\sqrt{2}} \gamma_\mu (1-\gamma_5)(\hat{P} - \hat{k} - \hat{l}) 2\pi \delta^+((P-k-l)^2) \\
& (-i) \frac{G_F}{\sqrt{2}} \gamma_\nu (1-\gamma_5) i \frac{-(\hat{k}+\hat{p}_4) + m'}{(k+p_4)^2 - m'^2 + i\epsilon} i g \gamma_\rho v_u(p_4) \\
& \frac{-i g^{\rho\sigma}}{k^2 - \lambda^2 + i\epsilon} \overline{u}_b(p_3) \gamma^\mu (1-\gamma_5) \hat{l} 2\pi \delta^+(l^2) \gamma^\nu (1-\gamma_5) u_b(p_1) \\
& (-2\pi i) \; \{ \; ((k+p_4)^2 - m'^2) \delta^+((k+p_4)^2 - m'^2) + \\
& (k^2 - \lambda^2) \delta^+(k^2 - \lambda^2) + ((k+p_2)^2 - m^2) \delta^+((k+p_2)^2 - m^2) \}
\end{aligned} \tag{8}$$

The gluon propagator is taken for simplicity in the Feynman gauge and infrared divergences are regulated with a fictious gluon mass λ. $P = p_1 - p_4$, p_1 and p_2 are the initial momenta of the b and $\overline{u}$ quark respectively, and p_3 and p_4 are the final momenta. The masses of the c and the d quarks have been taken equal to zero since they do not give rise to mass singularities (see later for the definition of m and m'). $C = C^{ik}_{je} = 1/2(C_+^2 + C_-^2)(1 - 1/N^2)\delta^i_j \delta^k_e$ is a color factor.

Computing the spin algebra, we see that the amplitude $\mathcal{M}$ can be decomposed according to the powers of the masses m and m' appearing at the numerator:

$$\mathcal{M} \;=\; \mathcal{M}^{(0)} + m' \mathcal{M}^{(1a)} + m \mathcal{M}^{(1b)} + m m' \mathcal{M}^{(2)} \tag{9}$$

The loop over l is easily integrated and we have:

$$\mathcal{M}^{(0)} \;=\; -\frac{\alpha_s G_F^2 C}{2\pi^3} \overline{v}_u(p_2) \gamma^\nu (1-\gamma_5) v_u(p_4) \overline{u}_b(p_3) \gamma^\mu (1-\gamma_5) u_b(p_1) I_{\mu\nu} \tag{10}$$

$$
\begin{aligned}
\mathcal{M}^{(1a)} &= -\frac{\alpha_s G_F^2 C}{\pi^3}\overline{v}_u(p_2)(1+\gamma_5)v_u(p_4)\overline{u}_b(p_3)\gamma^\mu(1-\gamma_5)u_b(p_1)V_\mu && (11)\\
\mathcal{M}^{(1b)} &= -\frac{\alpha_s G_F^2 C}{\pi^3}\overline{v}_u(p_2)(1-\gamma_5)v_u(p_4)\overline{u}_b(p_3)\gamma^\mu(1-\gamma_5)u_b(p_1)W_\mu && (12)\\
\mathcal{M}^{(2)} &= -\frac{\alpha_s G_F^2 C}{2\pi^3}\overline{v}_u(p_2)\gamma^\mu(1+\gamma_5)v_u(p_4)\overline{u}_b(p_3)\gamma_\mu(1-\gamma_5)u_b(p_1)S && (13)
\end{aligned}
$$

where $I_{\mu\nu} = I_{\mu\nu}^{(i)} + I_{\mu\nu}^{(ii)} + I_{\mu\nu}^{(iii)}$ and similarly for the other integrals. For the cut (iii), we have:

$$
\begin{aligned}
I_{\mu\nu}^{(iii)} &= \int \frac{d^4k\ \delta^+((k+p_2)^2-m^2)}{(k^2-\lambda^2+i\epsilon)((k+p_4)^2-m'^2+i\epsilon)}\{(k+p_2)_\mu(k+p_4)_\nu \qquad (14)\\
&\quad + (k+p_2)_\nu(k+p_4)_\mu - g_{\mu\nu}(k+p_2)(k+p_4)\\
&\quad - i\epsilon_{\rho\sigma\mu\nu}(k+p_2)^\rho(k+p_4)^\sigma\ \}\ \frac{1}{2}(P-k)^2\Theta^+((P-k)^2)\\
V_\mu^{(iii)} &= \int \frac{d^4k\ \delta^+((k+p_2)^2-m^2)}{(k^2-\lambda^2+i\epsilon)((k+p_4)^2-m'^2+i\epsilon)}(k+p_2)_\mu \times\\
&\qquad \times \frac{1}{2}(P-k)^2\Theta^+((P-k)^2)\\
W_\mu^{(iii)} &= \int \frac{d^4k\ \delta^+((k+p_2)^2-m^2)}{(k^2-\lambda^2+i\epsilon)((k+p_4)^2-m'^2+i\epsilon)}(k+p_4)_\mu \times\\
&\qquad \times \frac{1}{2}(P-k)^2\Theta^+((P-k)^2)\\
S^{(iii)} &= \int \frac{d^4k\ \delta^+((k+p_2)^2-m^2)}{(k^2-\lambda^2+i\epsilon)((k+p_4)^2-m'^2+i\epsilon)} \times\\
&\qquad \times \frac{1}{2}(P-k)^2\Theta^+((P-k)^2)
\end{aligned}
$$

where we have defined $\Theta^+(v^2) = \Theta(v^2)\Theta(v_0)$.
The corresponding integrals for the cuts (i) and (ii) are obtained from the cut (iii) respectively, with the substitutions:

$$
\begin{aligned}
\frac{\delta^+((k+p_2)^2-m^2)}{(k^2-\lambda^2)((k+p_4)^2-m'^2)} &\to \frac{\delta^+((k+p_4)^2-m'^2)}{(k^2-\lambda^2)((k+p_2)^2-m^2)} \qquad (15)\\
&\to \frac{\delta^+(k^2-\lambda^2)}{((k+p_2)^2-m^2)((k+p_4)^2-m'^2)}
\end{aligned}
$$

To simplify the computation of the integrals, we consider a non-relativistic configuration of the external momenta, i.e. we take:

$$p_1 = p_3 = (m_b, \vec{0}) \quad \text{and} \quad p_2 = p_4 = (m, \vec{0}) \tag{16}$$

The cuts (i) and (iii) have singularities in the forward limit, i.e. for $p_1 = p_3$ and $p_2 = p_4$, and the configuration (16) is a forward one. We regulate these singularities by taking a different mass for the $\overline{u}$ quark in the initial state, $m_u = m$, and in the final state, $m_u = m'$. After summing the cuts (i) and (iii), we are allowed to take the forward scattering limit $m' \to m$.
With our choice of the external momenta, $I_{\mu\nu}$ can be parametrized as:

$$I_{\mu\nu} \;=\; g_{\mu\nu} A + \frac{p_{2\mu} p_{2\nu}}{m^2} B \tag{17}$$

For the projections

$$X \;=\; g_{\mu\nu} I^{\mu\nu} \;=\; 4A + B \qquad \text{and} \qquad Y \;=\; \frac{p_{2\mu} p_{2\nu}}{m^2} I^{\mu\nu} \;=\; A + B, \tag{18}$$

we have, for the cut (iii):

$$\begin{aligned} X_{(iii)} \;&=\; \frac{-\pi}{\delta m} \int_0^{\overline{k}/m} \frac{k^2 dk}{2\sqrt{1+k^2}(\sqrt{1+k^2}-1)(\sqrt{1+k^2}-1+\lambda^2/2m^2)} \\ &\times\; [m - \delta m \sqrt{1+k^2}][(M+m)^2 + m^2 - 2m(M+m)\sqrt{1+k^2}] \\ &=\; \pi M^2 [\frac{\pi m}{\lambda} + \frac{1}{8}\frac{M}{m} - \frac{\pi m^2}{\delta m \lambda} + \ldots] \end{aligned} \tag{19}$$

and

$$\begin{aligned} Y_{(iii)} \;&=\; \frac{\pi}{\delta m} \int_0^{\overline{k}/m} \frac{k^2 dk}{2\sqrt{1+k^2}(\sqrt{1+k^2}-1)(\sqrt{1+k^2}-1+\lambda^2/2m^2)} \\ &\times\; \frac{1}{2}[(M+m)^2 + m^2 - 2m(M+m)\sqrt{1+k^2}][2mk^2 + m - \delta m\sqrt{1+k^2}] \\ &=\; \pi M^2 [-\frac{\pi}{2}\frac{m}{\lambda} - \frac{1}{16}\frac{M}{m} + \frac{1}{48}\frac{M^2}{m \delta m} + \frac{\pi}{2}\frac{m^2}{\delta m \lambda} + \ldots] \end{aligned} \tag{20}$$

In eqs.(19) and (20) the shift $k \to k + p_2$ and the conversion to an adimensional momentum k/m have been done.

The dots '…' indicate finite or at most logarithmic terms, $\delta m = m - m'$, $M = m_b - m$ is the temporal component of P, $P = (M, \vec{0})$, and $\overline{k} = \overline{k}(M, m)$ is the upper limit for the gluon 3-momenta:

$$\overline{k} = \sqrt{\left(\frac{M+m}{2}\right)^2 [1 + \left(\frac{m}{M+m}\right)^2] - m^2} = \frac{M}{2}[1 + \frac{m}{M} + \ldots] \tag{21}$$

The analyticity argument is forced to disregard the θ-function, which is taken into account in eq.(21).
$X_{(iii)}$ and $Y_{(iii)}$ contain power singularities of the form:

$$\frac{1}{\lambda}, \quad \frac{1}{\delta m} \quad \text{and} \quad \frac{1}{m} \tag{22}$$

$X_{(i)}$ and $Y_{(i)}$ are computed from $X_{(iii)}$ and $Y_{(iii)}$ respectively by exchanging m' with m. Summing the contributions from the two cuts, we see that singularities in $1/\lambda$ and $1/\delta m$ cancel, to give:

$$X_{(i)} + X_{(iii)} \;=\; \pi M^2 [\frac{1}{4}\frac{M}{m} + \ldots] \tag{23}$$

and

$$Y_{(i)} + Y_{(iii)} \;=\; \pi M^2 [-\frac{1}{48}\left(\frac{M}{m}\right)^2 - \frac{1}{8}\frac{M}{m} + \ldots] \tag{24}$$

The computation of the cut (ii) related to the real gluon emission diagram is simpler because there is no the forward singularity (one can take $m = m'$ from the beginning), and gives:

$$X_{(ii)} \;=\; \pi M^2 \frac{1}{4}[-\frac{M}{m} + \ldots] \tag{25}$$

and

$$Y_{(ii)} = \pi M^2 [\frac{1}{48}\left(\frac{M}{m}\right)^2 + \frac{1}{8}\frac{M}{m} + \ldots] \tag{26}$$

Summing the contributions from cuts (i), (ii) and (iii), we see that there is a cancellation of the power singularities in $1/m^2$ and $1/m$.

The computation of $V_\mu^{(iii)}$ gives:

$$V_\mu^{(iii)} = \delta_{\mu 0} \pi M^2 [\;\frac{\pi}{2}\frac{m}{\delta m \lambda} + \frac{1}{16}\frac{M}{m \delta m} + \ldots\;] \tag{27}$$

$V_\mu^{(i)}$ may be computed from $V_\mu^{(iii)}$ by means of the formula:

$$V_\mu^{(i)} \;=\; V_\mu^{(iii)}(m \to m') + \delta_{\mu 0}\;\delta m\; S^{(iii)}(m \to m') \tag{28}$$

The sum of the cuts (i) and (iii) gives in this case:

$$m[V_\mu^{(i)} + V_\mu^{(iii)}] \;=\; \delta_{\mu 0}\pi M^2[-\frac{1}{16}\frac{M}{m} + \ldots] \tag{29}$$

while

$$mV_\mu^{(ii)} \;=\; \delta_{\mu 0}\pi M^2[\;\frac{1}{16}\frac{M}{m} + \ldots] \tag{30}$$

The sum of the cuts does not contain any more power singularities. The computation of W_μ is completely analogous to that of V_μ and we do not report the results. For S we have the expression:

$$S_{(iii)} \;=\; \pi M^2[\;\frac{\pi}{2}\frac{1}{\delta m\lambda} + \frac{1}{4}\frac{1}{m\delta m}\log\frac{M}{m} + \ldots\;] \tag{31}$$

Using the symmetry relation $S_{(i)} = S_{(iii)}(m \to m')$, we have for the partial sum:

$$m^2[\;S_{(i)} + S_{(iii)}\;] = \pi M^2[-\frac{1}{4}\log\frac{M}{m} + \ldots] \tag{32}$$

Note that there is not any power singularity because of the m^2 at the numerator coming from Dirac algebra. $m^2 S_{(ii)}$ also is free from power singularities:

$$m^2 S_{(ii)} \;=\; \pi M^2[\;\frac{1}{4}\log\frac{M}{\lambda} + \ldots\;] \tag{33}$$

In the total sum therefore we have only an infrared logarithmic singularity:

$$m^2[\;S_{(i)} + S_{(ii)} + S_{(iii)}\;] \;=\; \pi M^2[\;\frac{1}{4}\log\frac{m}{\lambda} + \ldots\;] \tag{34}$$

This infrared singularity, together with those neglected in eqs.(23)-(30), cancels in the total sum of the real and virtual gluon emission diagrams.

We consider now the diagrams in figs.2 and 3, connecting an external heavy line with an external light line via a gluon. The cuts (i) on the gluon line have $1/m$ singularities, while the cuts (ii) on the light quark line have both $1/m$ and $1/\lambda$ singularities. $1/m$ singularities cancel in the sum of the cuts (i) and (ii) of each diagram separately, while $1/\lambda$ singularities cancel in the total sum of the four cuts. The computation is analogous to the one described and we do not give the details.

In conclusion, we have verified explicitly the cancellation of power enhancements in the inclusive widths of non-spectator decays.

Acknowledgment

I wish to express particular thanks to Profs. G. Altarelli and S. Petrarca for many useful discussions and for encouragement.

References

[1] J.L. Cortes, X.Y. Pham and A. Tounsi, Phys. Rev. D 25 (1982) 188.

[2] G. Altarelli and L. Maiani, Phys. Lett. B 52 (1974) 351; M.K. Gaillard and B.W. Lee, Phys. Rev. Lett. 33 (1974) 108.

[3] G. Altarelli et al., Phys. Lett. B 99 (1981) 141; Nucl. Phys. B 187 (1981) 461; A.J. Buras and H. Weisz, Nucl. Phys. B 333 (1990) 66.

[4] G. Altarelli and S. Petrarca, Phys. Lett. B 261 (1991) 303.

[5] Review of Particle Properties (1992).

[6] G. Altarelli et al., Nucl. Phys. B 208 (1982) 365.

[7] For a recent review on the Heavy Quark Effective Theory, with original references, M. Wise, Caltech preprint CALT-68-1901 (1993).

[8] I.I. Bigi, N.G. Uraltsev and A.I. Vainshtein, Phys. Lett. B 293 (1992) 430; erratum-ibidem B 297 (1993) 477; B. Blok and M. Shifman, NSF-ITP-92-103, TPI-MINN-92-33/T; TPI-MINN-92-52-T, NSF-ITP-92-115.

[9] M. Bander, D. Silverman and A. Soni, Phys. Rev. Lett. 44 (1980) 7; H. Fritzsch and P. Minkowski, Phys. Lett. B (1980) 455; Nucl. Phys. B 171 (1980) 413.

[10] For a recent determination of f_B with lattice QCD, see, for example, C.R. Allton et al., Rome preprint ROME 93/928.

[11] B. Guberina et al., Phys. Lett. B 89 (1979) 111; R.D. Peccei and R. Ruckl, Proc. Ahrenshoop Symp. (Ahrenshoop, German Democratic Republic, 1981); Max-Plank Institute, Munich preprint MPI-PAE/PTh 75/81 T. Kobayashi and N. Yamazaki, Prog. Theor. Phys. 65 (1981) 775; G. Altarelli and L. Maiani, Phys. Lett. B 118 (1982) 414.

[12] M.B. Voloshin and M.A. Shifman, Sov. J. Nucl. Phys. 41 (1985) 120; Sov. Phys. JETP 64 (1986) 698.

[13] M.B. Voloshin and M.A. Shifman, Yad. Fiz. 45 (1987) 463; H.D. Politzer and M. Wise, Phys. Lett. B 206 (1988) 681.

[14] I.I. Bigi and N.G. Uraltsev, Phys. Lett. B 280 (1992) 271.


$l, \ c$
b
p_1
$P-k-l, \ d$
p_3
$k \rightarrow$
$\overline{u}$
$p_2 \rightarrow$
$p_4 \rightarrow$
(i)
(ii)
(iii)

Fig. 1


(i)
(ii)

Fig. 2



Fig. 3